

AD-A248 026

2

Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release Distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  Technical Report No. 7			7a. NAME OF MONITORING ORGANIZATION  Office of Naval Research		
6a. NAME OF PERFORMING ORGANIZATION  Brigham Young University		6b. OFFICE SYMBOL (If applicable) BYU	7b. ADDRESS (City, State, and ZIP Code)  Department of the Navy Arlington, VA 22217-5000		
6c. ADDRESS (City, State, and ZIP Code)  Department of Chemistry Provo, UT 84602		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  N00014-91-J-1710			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION  Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR	10. SOURCE OF FUNDING NUMBERS		
8c. ADDRESS (City, State, and ZIP Code) 800 No. Quincy Street Arlington, VA 22217-5000		PROGRAM ELEMENT NO	PROJECT NO.	TASK NO	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Recognition of the Enantiomers of Chiral Organic Ammonium Salts By Chiral Dibenzyl- and Diphenyl-Substituted Diamido-or Dithionoamidopyridino-18-crown-6 Ligands					
12. PERSONAL AUTHOR(S) P. Huszthvy, J.S. Bradshaw, C.Y. Zhu, T. Wang, and R.M. Izatt					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1992, March 24	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>New chiral pyridino-18-crown-6 ligands containing two amide, thionoamide or amine units and two benzyl or phenyl substituents on chiral macroring positions have been synthesized (see Figure 1). Some of these chiral ligands exhibit only moderate or no recognition for the enantiomers of [<math>\alpha</math>-(1-naphthyl)ethyl]ammonium perchlorate as determined by <math>^1\text{H}</math> NMR spectral techniques.</p>					
CONTINUED ON BACK					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Harold Guard			22b. TELEPHONE (Include Area Code) (202) 696-4409		22c. OFFICE SYMBOL ONR

DD Form 1473, JUN 86

92-07983




3

SECURITY CLASSIFICATION OF THIS PAGE  
Unclassified

92 3 20 930

Figure 1. Chiral Diamido-, Dithionoamido-, Diaza- and Azapyridino-18-crown-6 Ligands




- |   |                                   |
|---|-----------------------------------|
| 1, X = NH; Y = O; R = benzyl (S,S)                              |                                   |
| 2, X = NH; Y = S; R = benzyl (S,S)                              |                                   |
| 3, X = NH; Y = H <sub>2</sub> ; R = benzyl (S,S)                |                                   |
| 4, X = NH; Y = O; R = phenyl (S,S)                              |                                   |
| 5, X = NH; Y = S; R = phenyl (S,S)                              |                                   |
| 6, X = NH; Y = H <sub>2</sub> ; R = phenyl (S,S)                |                                   |
| 7, X = NCH <sub>3</sub> ; Y = O; R = phenyl (S,S)               | 10, R = H (S,S)                   |
| 8, X = NCH <sub>3</sub> ; Y = S; R = phenyl (S,S)               | 11, R = C(O)CH <sub>3</sub> (S,S) |
| 9, X = NCH <sub>3</sub> ; Y = H <sub>2</sub> ; R = phenyl (S,S) |                                   |

Complexation of the enantiomeric forms of NapEt by some of the new ligands has been studied by the temperature-dependent <sup>1</sup>H NMR technique<sup>1-5</sup>, and by determining the log K values for the association of the chiral ligand and the enantiomer salt by a direct <sup>1</sup>H NMR spectral method.<sup>5,9</sup> Table I contains the free energy of activation ( $\Delta G_c^\ddagger$ ) and log K values for the interaction of some of these new chiral ligands with the enantiomers of [ $\alpha$ -(1-naphthyl)ethyl]ammonium perchlorate(NapEt). The data show that, in general, these chiral ligands exhibit only moderate or no recognition for the enantiomers of NapEt. The  $\Delta G_c^\ddagger$  data show that ligands 7, 8, and 9 form stable

Accession For	
Y-1000	<input checked="" type="checkbox"/>
Y-1000	<input type="checkbox"/>
Y-1000	<input type="checkbox"/>
Y-1000	<input type="checkbox"/>
By	
Distribution/	
Availability Codes	
Avail and/or	
Special	

A-1



complexes in  $\text{CD}_2\text{Cl}_2$  with the (R) and (S) forms of NapEt but the resulting complexes each have about the same kinetic stability. This is in sharp contrast to complexation by the chiral diphenyl-substituted pyridino-18-crown-6 ( $\text{Y} = \text{H}_2$ ,  $\text{X} = \text{O}$ ,  $\text{R} = \text{phenyl}$ ) and diesterpyridino-18-crown-6 ( $\text{Y} = \text{O}$ ,  $\text{X} = \text{O}$ ,  $\text{R} = \text{phenyl}$ ) ligands (see Figure 1) which exhibited considerable recognition of one form of NapEt over the other form in  $\text{CD}_2\text{Cl}_2$  as determined by the temperature dependent  $^1\text{H}$  NMR method.<sup>5</sup> Log K values in Table I do show that some of these ligands exhibit enantiomeric recognition in a mixture of 50%  $\text{CDCl}_3$  - 50%  $\text{CD}_3\text{OD}$  or in  $\text{CD}_3\text{NO}_2$ . Dithionoamide (S,S)-5 favored (R)-NapEt over (S)-NapEt by 0.37 log K units. This recognition compares favorably with that obtained by the similar diphenyl-substituted crown ( $\text{X} = \text{O}$ ,  $\text{Y} = \text{O}$ ) ligand.<sup>5</sup> Ligand (S,S)-8 also exhibited moderate recognition for (R)-NapEt in the 50%  $\text{CDCl}_3$  - 50%  $\text{CD}_3\text{OD}$  mixture. Ligands (S,S)-1 and (S,S)-7 exhibit some recognition for one form of NapEt in  $\text{CD}_3\text{NO}_2$ . Thus, in some cases, these new chiral ligands show moderate enantiomeric recognition but they have inferior recognition properties as compared to the diphenyl- and di-t-butyl-substituted pyridino- and diesterpyridino-18-crown-6 ligands (all have  $\text{X} = \text{O}$ ).<sup>5</sup>

The data in Table I also show that log K values for macrocycle-ammonium salt interactions increase as the polarity of the solvent decreases. This is an expected trend for solvent effects in these types of interactions. It is also instructive to note that there is a significant decrease in log K values when substituting an amide group for an ester in these systems. This is best observed by comparing the log K value of 0.7 (Table 1) for the interaction of (S,S)-4 with (R)-

NapEt in a mixture of 50%  $\text{CDCl}_3$  - 50%  $\text{CD}_3\text{OD}$  with a log  $K$  value of 2.15 for the same interaction of the comparable ester ( $X$  and  $Y = \text{O}$ ,  $R = \text{phenyl}$ ) in a mixture of 30%  $\text{CDCl}_3$  - 70%  $\text{CD}_3\text{OD}$ .<sup>5</sup> The two solvent systems are slightly different but the latter solvent is more polar which should decrease the log  $K$  value. This significant decrease in complexing ability for the diamido-crowns is probably a result of distorted macrorings that will be mentioned later. The diamino-crowns, on the other hand, form stronger complexes as shown by the log  $K$  value of 3.2 for the interaction of ( $S,S$ )-9 with ( $R$ )-NapEt.

Table I. Free Energies of Activation ( $\Delta G_c^\ddagger$ )<sup>a</sup> and log K Values<sup>b</sup> for the Interaction of Some Chiral Macrocyclic Ligands with the R and S Forms of [ $\alpha$ -(1-Naphthyl)ethyl]ammonium Perchlorate (NapEt)

Ligand	$\Delta G_c^\ddagger$ values (Kcal/mol) <sup>c</sup>		log K values at 20 °C		
	( <u>R</u> )-NapEt	( <u>S</u> )-NapEt	( <u>R</u> )-NapEt <sup>d</sup>	( <u>S</u> )-NapEt <sup>d</sup>	( <u>S</u> )-NapEt <sup>e</sup>
( <u>S</u> , <u>S</u> )-1			g	g	0.8
( <u>S</u> , <u>S</u> )-4	f	f	0.7	g	
( <u>S</u> , <u>S</u> )-5			1.39	1.02	
( <u>S</u> , <u>S</u> )-7	11.3	11.2	g	g	2.75
( <u>S</u> , <u>S</u> )-8	>14.0	>13.9	0.7	0.5	
( <u>S</u> , <u>S</u> )-9	12.0	12.1	3.2	3.3	h

<sup>a</sup> A Varian VXR-500 spectrometer was used to record all <sup>1</sup>H NMR spectra for determination of both  $\Delta G_c^\ddagger$  and log K values. The procedure for the determination of  $\Delta G_c^\ddagger$  values was as described in ref. 1-5.

<sup>b</sup> Log K values were determined by a direct <sup>1</sup>H NMR procedure as described in ref. 9.

<sup>c</sup> CD<sub>2</sub>Cl<sub>2</sub> was used as a solvent.

<sup>d</sup> 50%/50% (v/v) CDCl<sub>3</sub>-CD<sub>3</sub>OD mixture was used as a solvent.

<sup>e</sup> CD<sub>3</sub>NO<sub>2</sub> was used as a solvent.

<sup>f</sup> No splitting was observed in the <sup>1</sup>H NMR spectra.

<sup>g</sup> Log K value was so low that accurate measurements could not be made.

<sup>h</sup> Log K value was so high that accurate calculations could not be made.

## References

- (1) Davidson, R. B.; Bradshaw, J. S.; Jones, B. A.; Dalley, N. K.; Christensen, J. J.; Izatt, R. M.; Morin, F. G.; Grant, D. M. J. Org. Chem. 1984, 49, 353.
- (2) Bradshaw, J. S.; Thompson, P. K.; Izatt, R. M.; Morin, F. G.; Grant, D. M. J. Heterocyclic Chem. 1984, 21, 897.
- (3) Bradshaw, J. S.; Colter, M. L.; Nakatsuji, Y.; Spencer, N. O.; Brown, M. F.; Izatt, R. M.; Arena, G.; Tse, P.-K.; Wilson, B. E.; Lamb, J. D.; Dalley, N. K.; Morin, F. G.; Grant, D. M. J. Org. Chem. 1985, 50, 4865.
- (4) Bradshaw, J. S.; Huszthy, P.; McDaniel, C. W.; Zhu, C. Y.; Dalley, N. K.; Izatt, R. M.; Lifson, S. J. Org. Chem. 1990, 55, 3129.
- (5) Huszthy, P.; Bradshaw, J. S.; Zhu, C. Y.; Izatt, R. M.; Lifson, S. J. Org. Chem. 1991, 56, 3330.
- (6) Kyba, E. B.; Koga, K.; Sousa, L. R.; Siegel, M. G.; Cram, D. S. J. Am. Chem. Soc. 1973, 95, 2692.
- (7) Li, Y.; Echegoyen, L.; Martinez-Diaz, M. V.; Mendoza, J.; Torres, T. J. Org. Chem. 1991, 56, 4193.
- (8) Izatt, R. M.; Zhu, C.-Y.; Huszthy, P.; Bradshaw, J. S.  
"Enantiomeric Recognition in Macrocycle-Primary Ammonium Cation Systems," in "Crown Ethers: Toward Future Applications," Cooper, S. R. Ed., VCH Press, New York, in press.
- (9) Zhu, C. Y.; Bradshaw, J. S.; Oscarson, S. L.; Izatt, R. M. J. Incl. Phenom. in press.